REFERENCES

1)The Merck Index, Ed.S.Budavari, (2006), Susan, Merck Research Lab., White station Division of Merck &co., Inc. 14th. Edition, 1569**.**

2)Hillaert S,(2003), *J Pharm Biomed Anal*, **31,** 329.

3)Sethi P.D, (1996), High Performance Thin Layer Chromatography, Quantative Analysis of Pharmaceutical Formulations. CBS Publishers and Distributors, New Delhi.

4)Harpey C, Clauser P, Labird C, Freyria J, and Poirier J, (1989), Cardiovascular Drugs Reviews, **6(4),**292-312.

5)Physician Desk Reference, (2000), 54th edition, Medical Economic Company, 2980- 84.

6)Gorge S, (1997) ,The Essential Guide to Analytical Chemistry, John Willy and Sons.

7)Chen, Ruichun, Jia, Haitao, Ai, Lianfeng; Guo, Chunha, (2011), Determination of Nine Penicillin Residues in Milk and Milk Powder by HPLC-MS/MS, *shipin kexue* **32(18)**, 249-252

8)Samanidou, Victoria F, Giannakis, Dimitrios E, Papadaki, Antonia, (2009), Development and validation of an HPLC method for the determination of seven penicillin antibiotics in veterinary drugs and bovine blood plasma**,***Journal of separation science* **32(9)**, 1302-1311

9)Hong, Jianwen; Wei, Xuefang; Yu, Caidi, (2006), Comparative Study of RP-HPLC and UV Spectrophotometric Techniques for the Simultaneous Determination of Amoxicillin and Cloxacillin in Capsules,Guangdong Yaoxueyuan Xuebao, 22(6), 624-626

10)Zhang, Mengjun, Zhang, Huijing, Guan, Xiao,Xiao, Zhenghua, (2006), Development and Validation of Reversed-PhaseHPLC Method for Simultaneous Estimation of Rosuvastatin and Fenofibrate in Tablet Dosage Form,*Yaowu Fenxi Zazhi*, **26(2)**, 228-231

11)Sorensen, L. K. Snor, L. K,(2001), *Chromatogaphia, Determination of ampoxillin by Hplc*, **53(7/8)**, 367-371

12)El Sayed, Khaled S, Younis, Mohamed E. El-Shanawany, Abdalla A. Zagazig (1998), A simple HPLC procedure was described for the detn. of amoxicillin, ampicillin, cloxacillin, dicloxacillin, flucloxacillin andphenoxymethylpenicillin (penicillin V) in single or combined pharmaceutical formulations,*Journal of Pharmaceutical sciences*, **7(1)**, 38-47.

13)Hong, Chih-Chun, Lin, Chin-Lan, Tsai, Chin-En; Kondo, Fusao,(1995), A new improved RP-HPLC method for assay of rosuvastatin calcium in tablets*American Journal of Research* **56(3)**, 297-303**.**

14)Devika, G. S, Sudhakar M, Venkateshwara Rao, (2011), A new improved RP-HPLC method for assay of rosuvastatin calcium in tablets, *International Journal of Pharmacy and Pharmaceutical Sciences* , 311-315

15)Sevda, R. R, Ravetkar A. S, Shirote P. J,(2011), Spectrophotometric estimation of Rosuvastatin. Calcium by UV method,*International Journal of ChemTech Research,* **3(2)**, 629-635

16)Wang, Tiesong, Tong, Yu, Zheng, Jie, Zhang, Zhe, Gao, Qing, Che, Baoquan, (2010), Detection of 29 Adulterations in Lipid-Regulating, Antihypertensive, Antitussive and Antiasthmatic Herbal Remedies by TLC *Zhongguo Yaoxue Zazhi*, **45(11)**, 857-861

17)Thimmaraju, Manish Kumar, Rao, Venkat, Gurrala, Srikanth,(2011), RP HPLC method for the determination of finasteride and tamsulosin in bulk and pharmaceutical formulations, *pharmacia Lettre,* **3(5)**, 79-86

18)Thimmaraju, Manish Kumar, Rao, Venkat, Gurrala, Srikanth, Reddy, Jayapal G, (2011), Detection of 29 Adulterations in Lipid-Regulating, Antihypertensive, Antitussive and Antiasthmatic Herbal Remedies by TLC *International Journal of pharmacy and biological sciences*, **1(3)**, 303-310

19)Gadhave, Nilam A, Ghante, Minal R, Nikam, Atul D, Sawant, Sanjay D, (2011), Simultaneous estimation of tamsulosin hydrochloride and finasteride in combined dosage form by UV spectroscopy method,*Journal of Pharmacy Research,* **4(8)**, 2672-2674

20)Bari, Sanjay B, Jain, Pritam S. Bakshi, Abdul R, Surana, Sanjay J, (2011), HPTLC Method Validation for simultaneous determination of Tamsulosin Hydrochloride and Finasteride in Bulk and Pharmaceutical Dosage Form, *Journal of Analytical & Bioanalytical Techniques*, **2(2)**, 1000119

21)Kategaonkar, Atul H, Patel, Dhaval M. Itkar, Bhushan P, Choudhari, Vishnu P, Kuchekar, Bhanudas S, Nikalje, Ana G, (2009), Spectrophotometric and Stability Indicating

RP-HPLC-PDA method for Simultaneous Determination of Finasteride and Tamsulosin in Combined Tablet Dosage form, *Asian Journal of Research in Chemistry,* **2(4)**, 414-419

22)Patel, Dipti B, Patel, Natubhai J, (2010), Validated RP-HPLC and TLC methods for simultaneousestimation of tamsulosin hydrochloride and finasteride in combined dosage forms*Acta Pharmaceutica* ,**60(2)**, 197-205

23)Wang, Dongkai, Qiu, Lipeng, Li, Xiangm, (2010), The title compound sustained-release capsule comprises: tamsulosin 0.01-10%, finasteride 0.1-20%, sustained-releaseadjuvant 60-97%, and other adjuvant 0-10%. Tamsulosin can be hydrochloride, hydroiodide, hydrobromide and fumarate of tamsulosin, Faming Zhuanli Shenqing,CN 101703512 A 20100512

24)Vaidya V.V, Roy, Shikha M. N, Mangaokar K. V, Kekare M. B, Choukekar M. P, (2008), A simple, fast and precise reversed phase high performance liquid chromatography method is developed for the simultaneous determination of diloxanide furoate and tinidazole using metronidazole as an internal standard, *Analytical Chemistry: An Indian Journal*,**7(7)**, 505-508

25)Argekar, A. P, Powar S. G, Lemke, Thomas L, Williams, David A ,(1999), A simple, precise, rapid, and selective high-performance thin-layer chromatography (HPTLC) method was developed for the simultaneous determination of diloxanide furoate and tinidazole in tablets, *Journal of Planar Chromatography--Modern TLC,* **12(6)**, 452-455

26)Kale A. D, Kachhwaha S. J, Jadhav G. P, Trivedi, Piyush,

Lemke, Thomas L, Williams, David A,(1998), Dicyclomine was determined in the presence of tinidazole and diloxanide furoate by treatment with pH 1.2 potassiumdichromate solution and measurement of the absorbances at 357 nm. Beer's law held for 0-400 mg/mL, *Indian Drugs*, **35(11)**, 718-720

27)Podder G, Bandyopadhyay A, Chattopadhyay R. R, Maitra S. K, Ray S, Lemke, Thomas L, Williams, David A, (1989), Tinidazole (I) and diloxanide furoate (II) were determined in combined pharmaceuticals by using a double beam high-speed scanning spectrophotometer and measuring I and II at 310 and 254 nm, *Journal of the Institution of Chemists,* **61(5)**, 181-2

28)Ray S,Lemke, Thomas L, Williams, David A, (1989), A HPLC method for the simultaneous estimation of tinidazole and diloxanide furoate uses a micro Bondapak C18 column,MeOH-

0.05M H3PO4 (8:3) mobile phase and detection at 254 nm, *Eastern Pharmacist*, **32(381)**, 125-7

29)Sethi P. D, Chatterjee P. K, Jain C. L, Lemke, Thomas L. Williams, David A, (1988), Mixed dyslipidemia, oxidative stress and inflammation are related to a high risk for cardiovascular events ,*Journal of Pharmaceutical and Biomedical Analysis*, **6(3)**, 253-8

30)Pai P N, Rao G K, Srinivas B, Puranik S, (2008), A new improved RP-HPLC method for assay of rosuvastatin calcium in tablets*Indian journal of pharmaceutical sciences* **70(5)**, 670-2

31) Babu B, Hari, Rao M, Nageswara, Rambabu A, Srinivasan, P, (2011), A new simple, precise, economically viable and efficient High Performance Liquid Chromatography. (HPLC) method for the estimation of Methylcobalamin (MeB12) in bulk drugs and pharmaceutical dosage formulations was developed and validated, *Journal of Pharmacy Research*, **4(6)**,1685-1687

32)Agouridis, Aris P, Tsimihodimos, Vasilis, Filippatos, Theodosios D, Dimitriou, Andromachi A, Tellis, CostantinosC, Elisaf, Moses S, Mikhailidis, Dimitri P, Tselepis, Alexandros D,(2011), The effects of rosuvastatin alone or in combination with fenofibrate or omega 3 fatty acids on inflammation and oxidative stress in patients with mixed dyslipidemia,*Expert Opinion on Pharmacotherapy*, 12(17), 2605-2611

33)Darwish, Ibrahim A, Al-Obaid, Abdul-Rahman M, Al-Malaq, Hamoud A,(2011), Generation of polyclonal antibody with high avidity to rosuvastatin and its use in development of highly sensitive ELISA for determination of rosuvastatin in plasma *Chemistry Central Journal*, **5**, 38

34)*Faming Zhuanli Shenqing*, (2011), The invention relates to a synergistic pharmaceutical formulation for preventing and/or treating cardiovascular andcerebrovascular diseases, CN 101985042 A 20110316

35)Strain, Joe D, Farver, Debra K, Clem, James R,(2010), A review on the rationale and clinical use of concomitant rosuvastatin and fenofibrate/fenofibric acid therapy *Clinical Pharmacology: Advances and Applications*, **2**, 95-104

36)Li, Baoqi, Wang, Xiaodan, (2011), The title componud contains chemical combination of lipid-lowering drugs fenofibrate and nitric oxide donor drugs and onekind of HMG-CoA

reductase inhibitor as active ingredient, and pharmaceutically acceptable adjuvants,*Faming Zhuanli Shenqing* CN 101961490 A 20110202

37)Kida, Yasuo, Hasegawa, Masaaki, Kishi, Akio, Fukuda, Takuya, Joko, Mari, Shikano, Tsutomu, Kashiwagi, Atsunori, (2010), Synthesis of glycerolipid carbamates and dicarbamates and their use as an antitumor compounds,*Diabetes Frontier*, **21(5)**, 599-604

38)Yao, Yonggan, (2010), invention relates to choline salt of the medicine for lowering blood lipid, its preparaion and medical application ,*Faming Zhuanli Shenqing*, CN 101863780 A 20101020

39)Wang, Tiesong, Tong, Yu, Zheng, Jie, Zhang, Zhe, Gao, Qing, Che Baoquan, (2010), TLC methods for fast detection of lipid-regulating agents, antihypertensive agents, anti-tussive and antiasthmaticagents in herbal remedies simultaneously were established, *Zhongguo Yaoxue Zazhi*, **45(11)**, 857-861